

Table 2. Clinical outcomes of management of placenta accreta spectrum

| Variable | Uterus preserving management (n = 592, 81.43%) | Cesarean hysterectomy (n = 135, 18.56%) | p value |
|------------------------------------------------------|------------------------------------------------------|--------------------------------------------|----------|
| Intraoperative complications | | | < 0.0001 |
| <i>None</i> | 544 (91.89%) | 99 (73.33%) | 0.0001 |
| <i>Unintentional cystotomy</i> | 41 (6.93%) | 33 (24.44%) | < 0.0001 |
| <i>Ureteric injury</i> | 3 (0.51%) | 2 (1.48%) | 0.216 |
| <i>bowel injury</i> | 4 (0.68%) | 1 (0.74 %) | 0.934 |
| Cord clamping | | | > 0.0001 |
| <i>Immediate</i> | 511 (86.32%) | 95 (70.37%) | |
| <i>Delayed > 30 seconds</i> | 81 (13.68%) | 40 (29.63%) | |
| Estimated blood loss | 1200 (750 - 2000) | 1300 (750 – 2400) | 0.3021 |
| Massive blood loss (> 2500 ml) | 86 (14.5%) | 43 (31.9%) | < 0.0001 |
| DIC | 27 (4.56%) | 15 (11.11%) | 0.003 |
| Number of transfused units of packed RBCs | 0 (0-3) | 3 (2-4) | < 0.0001 |
| Number of transfused units of FFP | 0 (0-2) | 0 (0-3) | < 0.0001 |
| Maternal admission to ICU | 146 (24.66%) | 43 (31.85%) | 0.086 |
| Hospital stay in days | 4 (3-6) | 6 (4-8) | < 0.0001 |
| Postoperative hemoglobin (g/dl) | 9.5 (8.4- 10.6) | 8.9 (8- 10.1) | 0.0013 |
| IUFD | 11 (1.86%) | 0 (0%) | 0.111 |
| Postoperative AKI | 3 (0.51%) | 1 (0.74%) | 0.744 |
| Postoperative pulmonary oedema | 2 (0.34%) | 2 (1.48%) | 0.106 |

BMI: body mass index; CS: cesarean section; IVF: in-vitro fertilization; D & C: dilation and curettage; PAS: placenta accreta spectrum; DIC: disseminated intravascular coagulopathy; RBC: red blood cell; FFP: fresh frozen plasma; ICU: Intensive care unit