

Investigation of Exact Solutions of M-fractional Ivancevic Option Pricing Model Based on Three Different Methods

M. Raheel^{1,3*}, Khalid K. Ali^{2†}, Asim Zafar^{3‡}, Ahmet Bekir^{3§}

¹ Department of Mathematics & Statistics, Institute of Southern Punjab, Multan, Pakistan.

² Mathematics Department, Faculty of Science, Al-Azhar University, Nasr-City, Cairo, Egypt.

³Department of Mathematics, CUI Vehari Campus, Pakistan.

⁴Neighbourhood of Akcaglan, Imarli Street, Number: 28/4, 26030, Eskisehir, Turkey.

Abstract

This paper is about the investigation of exact solutions of important economic model; Ivancevic option pricing model (IOPM) with M-fractional derivative. To achieve this aim, three different methods; exp_a function method, extended Sinh-Gordon equation expansion method (EShGEEM) and extended (G'/G) -expansion method are used. Obtained solutions consisting of trigonometric, hyperbolic trigonometric, rational and exponential. The obtained solutions are new than the existing solutions in the literature. The got solutions are also verified by using Mathematica tool. Graphically justification are also done by plotting 2-D,3-D and contour graphs. The importance of this paper is that M-fractional derivative is first time use for this model. On the bases of achieved results it is suggested that these methods are simple, reliable and fruitful than the other methods.

Keywords: Ivancevic option pricing model; M-fractional derivative; exp_a function method; EShGEEM; Extended (G'/G) -expansion method; Exact solutions.

1 Introduction

Many mathematical models have been developed in these areas in the form of nonlinear partial differential equations (NLPDEs). Numerous schemes are made to gain exact solutions of NLPDEs like; generalized exponential rational function scheme (GERFS) [1–3],

*Corresponding author: mraheelmphil.math@gmail.com

†Corresponding author: khalidkaram2012@azhar.edu.eg

‡Corresponding author: asimzafar@hotmail.com

§Corresponding author: bekirahmet@gmail.com

Liu's extended trial function method (LETFM) [4], generalized unified method (GUM) [5], sine-Gordon expansion technique (SGET) [6], enhanced modified simple equation scheme [7], unified technique [8], extended tanh function scheme [9], Lie symmetry technique [10], symbolic computational method, Hirota's simple method and long wave technique [11], Jacobi elliptic function expansion scheme [12], Elzaki transform decomposition technique [13], $(m + \frac{1}{G})$ -expansion and adomian decomposition schemes [14], new generalized expansion method [15], simplest equation and kudryashov's new function techniques [16], modified simple equation scheme [17], modified kudryashov simple equation technique [18], first integral technique [19], Bäcklund transformation scheme [20], extended jacobi elliptic function expansion technique [21], extended (G/G) -expansion and improved (G'/G) -expansion schemes [30] etc.

There are three other methods; exp_a function, extended Sinh-Gordon equation expansion (EShGEE) and extended (G'/G) -expansion methods. These methods have various applications. Likely, some new kind of analytical results of perturbed Gerdjikov-Ivanov model (pGIM) has been achieved by using exp_a function and extended tanh function expansion methods in [31]. By applying exp_a function and hyperbolic function methods, various types of wave solutions of two non-linear Schrödinger equations are gained in [32]. New trigonometry and hyperbolic function type soliton solutions of (2+1)-dimensional hyperbolic and cubic-quintic non-linear Schrödinger equations are achieved by applying extended sinh-Gordon equation expansion scheme in [33]. Bright, dark and bright-dark soliton solutions of generalized non-linear Schrödinger equation has been determined by implementing extended sinh-Gordon equation expansion method in [34]. Some exact solitons of (2+1)-dimensional improved Eckhaus equation have been calculated by using extended (G'/G) -expansion technique in [35]. Various kinds of wave solutions of time-fractional parabolic equations have been obtained by applying the extended (G'/G) -expansion scheme in [36]. Fractional calculus have gained much importance due to its various applications in different fields. Therefore, different definition of derivatives have been used like; conformable fractional derivative [37,38], beta derivative [39], caputo-Fabrizio fractional derivative [40], truncated M-fractional derivative [41,42] etc.

Our considering model is one of the important and interesting economical model named as; Ivancevic option pricing model (IOPM). In the literature, few techniques have been used on this model to get different exact solutions. For example, new solutions have been achieved of this model by applying the fractional reduced differential transform technique in [43]. Dark, bright, dark-bright, complex, travelling, periodic, trigonometric, hyperbolic function solutions have been gained with the help of rational sine-Gordon expansion method and modified exponential method in [44]. Rogue wave and dark wave solitons of Ivancevic option pricing equation have been obtained with the use of trial function technique in [45]. The basic focus of the work is to investigate exact wave solutions of truncated M-fractional Ivancevic option pricing model based on exp_a function method, extended Sinh-Gordon equation expansion and extended (G'/G) -Expansion methods.

Paper have different sections; Section 2: truncated M-fractional derivative and its charac-

teristics, Section 3: model description, Section 4: description of methodologies, Section 5: mathematical treatment of model and exact solutions of model. Section 6: Illustrations with graphics. Section 7: conclusion.

2 Truncated M-derivative:

2.1 Definition:

Suppose $u(t) : [0, \infty) \rightarrow \mathbb{R}$, then truncated M-derivative of u of order ϵ is given [22]:

$$D_{M,t}^{\epsilon,\varrho} u(t) = \lim_{\tau \rightarrow 0} \frac{u(t E_{\varrho}(\tau t^{1-\epsilon})) - u(t)}{\tau}, \quad 0 < \epsilon < 1, \quad \varrho > 0, \quad (1)$$

where $E_{\varrho}(\cdot)$ shows truncated Mittag-Leffler function of one parameter that is defined as [23]:

$$E_{\varrho}(z) = \sum_{j=0}^{\infty} \frac{z^j}{\Gamma(\varrho j + 1)}, \quad \varrho > 0 \quad \text{and} \quad z \in \mathbf{C}. \quad (2)$$

2.2 Characteristics

Let $\epsilon \in (0, 1]$, $\varrho > 0$, $r, s \in \mathbb{R}$, and g, f ϵ -differentiable at a point $t > 0$, then by [22]:

$$(i) \quad D_{M,t}^{\epsilon,\varrho}(rg(t) + sf(t)) = rD_{M,t}^{\epsilon,\varrho}g(t) + sD_{M,t}^{\epsilon,\varrho}f(t). \quad (3)$$

$$(ii) \quad D_{M,t}^{\epsilon,\varrho}(g(t) \cdot f(t)) = g(t)D_{M,t}^{\epsilon,\varrho}f(t) + f(t)D_{M,t}^{\epsilon,\varrho}g(t). \quad (4)$$

$$(iii) \quad D_{M,t}^{\epsilon,\varrho}\left(\frac{g(t)}{f(t)}\right) = \frac{f(t)D_{M,t}^{\epsilon,\varrho}g(t) - g(t)D_{M,t}^{\epsilon,\varrho}f(t)}{(f(t))^2}. \quad (5)$$

$$(iv) \quad D_{M,t}^{\epsilon,\varrho}(A) = 0, \quad \text{where } A \text{ is a constant.} \quad (6)$$

$$(v) \quad D_{M,t}^{\epsilon,\varrho}g(t) = \frac{t^{1-\epsilon}}{\Gamma(\varrho + 1)} \frac{dg(t)}{dt}. \quad (7)$$

3 Model Description:

Let's assume M-fractional Ivancevic option pricing model(IOPM) [45] given as follows:

$$\iota D_{M,t}^{\epsilon,\varrho}q + \frac{\delta}{2} D_{M,2s}^{2\epsilon,\varrho}q + \Omega q|q|^2 = 0, \quad \iota = \sqrt{-1}. \quad (8)$$

This model was first time developed by Ivancevic [24] to fulfill both behavioral and efficient markets. Where $q = q(s, t)$ shows the option price wave profile. While t is the time variable and s is asset price of model. Parameter δ represents the volatility which shows either

stochastic process itself or only a constant. Where $\Omega = \Omega(r, \omega)$ is called Landau coefficient which describes adaptive market potential. In nonadaptive simplest case Ω and r become equal which shows the interest rate while in adaptive case $\Omega(r, \omega)$ may be connected to market temperature and it depends on the set of tractable parameters $\{W_i\}$. In third term $|q|^2$ shows the probability density function which denotes the potential field.

3.1 Summary of exp_a function scheme:

Here, we will give complete concept of this scheme.

Assuming the non-linear PDE;

$$G(q, q^2 q_t, q_x, q_{tt}, q_{xx}, q_{xt}, \dots) = 0. \quad (9)$$

Eq.(9) transformed in non-linear partial differential equation:

$$\Lambda(Q, Q', Q'', \dots) = 0. \quad (10)$$

By using following transformations:

$$q(x, y, t) = Q(\zeta), \zeta = ax + by + rt. \quad (11)$$

Considering root of Eq. (10) is shown in [25–28]:

$$Q(\zeta) = \frac{\alpha_0 + \alpha_1 d^\zeta + \dots + \alpha_m d^{m\zeta}}{\beta_0 + \beta_1 d^\zeta + \dots + \beta_m d^{m\zeta}}, \quad d \neq 0, 1. \quad (12)$$

where α_i and β_i ($0 \leq i \leq m$) are undetermined. Positive integral value of m is calculate by utilizing homogeneous balance technique into Eq.(10). Putting Eq. (12) into Eq. (10), gives

$$\wp(d^\zeta) = \ell_0 + \ell_1 d^\zeta + \dots + \ell_t d^{t\zeta} = 0. \quad (13)$$

Taking ℓ_i ($0 \leq i \leq t$) in Eq. (13) equal to 0, a system of algebraic equations is achieved as fellows.

$$\ell_i = 0, \quad \text{where } i = 0, \dots, t. \quad (14)$$

By using the got solutions, we achieve the exact results of Eq.(9).

3.2 Presentation of EShGEEM:

In this part, there are some fundamental steps of this method:

Step 1:

Assuming the NLPDE shown as follow:

$$G(q, D_{M,t}^{\alpha,\gamma} q, q^2 q_x, q_y, q_{yy}, q_{xx}, q_{xy}, \dots) = 0, \quad (15)$$

where $q = q(x, y, t)$ represents wave function.

Supposing the wave transform given as follow:

$$q(x, y, t) = Q(\zeta), \quad \zeta = x - \nu y + \frac{\Gamma(\gamma + 1)}{\alpha}(\kappa t^\alpha). \quad (16)$$

Inserting Eq. (16) into Eq. (15), resulting ordinary differential equation shown as follows:

$$\Lambda(Q(\zeta), Q^2(\zeta)Q'(\zeta), Q''(\zeta), \dots) = 0. \quad (17)$$

Step 2:

Let's assume the root of Eq. (16) in the form:

$$Q(p) = \alpha_0 + \sum_{i=1}^m (\beta_i \sinh(p) + \alpha_i \cosh(p))^i, \quad (18)$$

where $\alpha_0, \alpha_i, \beta_i$ ($i = 1, 2, 3, \dots, m$) are undetermined. A new function p of ζ that fulfill the following equation:

$$\frac{dp}{d\zeta} = \sinh(w). \quad (19)$$

Positive integral value of m may be obtained with the use of homogenous balance method. Eq. (19) is achieved from sinh-Gordon equation given as follows:

$$q_{xt} = \kappa \sinh(v). \quad (20)$$

Resultantly given in [29], one may get the roots of Eq. (20) shown as:

$$\sinh p(\zeta) = \pm \operatorname{csch}(\zeta) \quad \text{or} \quad \cosh p(\zeta) = \pm \operatorname{coth}(\zeta), \quad (21)$$

and

$$\sinh p(\zeta) = \pm \iota \operatorname{sech}(\zeta) \quad \text{or} \quad \cosh p(\zeta) = \pm \tanh(\zeta), \quad (22)$$

where $\iota = \sqrt{-1}$.

Step 3:

Substituting Eq. (18) along Eq. (20) into Eq. (17), result in the form of algebraic expressions in $p'^k(\zeta) \sinh^l p(\zeta) \cosh^m p(\zeta)$ ($k = 0, 1; \quad l = 0, 1; \quad m = 0, 1, 2, \dots$). Putting each coefficient of $p'^k(\zeta) \sinh^l p(\zeta) \cosh^m p(\zeta)$ equal to 0, to obtain a set of algebraic equations having $\nu, \kappa, \alpha_0, \alpha_i$ and β_i ($i = 1, 2, 3, \dots, m$).

Step 4:

By manipulating the gained set of algebraic equations by using Mathematica software, one may get the results of unknowns, $\nu, \kappa, \alpha_0, \alpha_i$ and β_i .

Step 5:

With the help of obtained solutions and Eqs. (21) and (22), we may achieve the roots of Eq. (17) shown as:

$$Q(\zeta) = \alpha_0 + \sum_{i=1}^m (\pm \beta_i \operatorname{csch}(\zeta) \pm \alpha_i \operatorname{coth}(\zeta))^i. \quad (23)$$

and

$$Q(\zeta) = \alpha_0 + \sum_{i=1}^m (\pm \beta_i \operatorname{sech}(\zeta) \pm \alpha_i \tanh(\zeta))^i. \quad (24)$$

3.3 Presentation of Extended (G'/G) -expansion Method:

In this part, there are some fundamental steps of this method given in [30].

Step 1: Supposing the NLPDE shown as follows:

$$G(q, D_{M,t}^{\alpha,\gamma} q, q^2 q_x, q_y, q_{yy}, q_{xx}, q_{xy}, \dots) = 0, \quad (25)$$

where $q = q(x, y, t)$ show the wave function.

Assuming the wave transform shown as follows:

Step 2:

$$q(x, y, t) = Q(\zeta), \quad \zeta = x - \nu y + \frac{\Gamma(\gamma + 1)}{\alpha} (\kappa t^\alpha). \quad (26)$$

Putting Eq. (26) into Eq. (25), results in the form of ODE shown as:

$$\Lambda(Q(\zeta), Q^2(\zeta)Q'(\zeta), Q''(\zeta), \dots) = 0. \quad (27)$$

Step 3:

Considering toots of Eq. (27) in the form given as:

$$Q(\zeta) = \sum_{i=-m}^m \alpha_i \left(\frac{G'(\zeta)}{G(\zeta)} \right)^i. \quad (28)$$

In Eq. (28), α_0 and α_i , ($i = \pm 1, \pm 2, \pm 3, \dots, \pm m$) are unknowns and $\alpha_i \neq 0$. Using homogenous balance method into Eq. (27), one can calculate positive integer m . Function $G = G(\zeta)$ fulfill the Riccati differential equation shown as follows:

$$dGG'' - aG^2 - bGG' - c(G')^2 = 0, \quad (29)$$

where a, b, c and d are constants.

Step 4:

Suppose Eq. (29) have results shown as:

Case 1: if $b \neq 0$ and $b^2 + 4ad - 4ac > 0$, then

$$\begin{aligned} \left(\frac{G'(\zeta)}{G(\zeta)} \right) &= \frac{b}{2(d-c)} \\ &+ \frac{\sqrt{-4ac + 4ad + b^2}}{2(d-c)} \left(\frac{C_1 \sinh \left(\frac{\zeta \sqrt{-4ac + 4ad + b^2}}{2d} \right) + C_2 \cosh \left(\frac{\zeta \sqrt{-4ac + 4ad + b^2}}{2d} \right)}{C_1 \cosh \left(\frac{\zeta \sqrt{-4ac + 4ad + b^2}}{2d} \right) + C_2 \sinh \left(\frac{\zeta \sqrt{-4ac + 4ad + b^2}}{2d} \right)} \right). \end{aligned} \quad (30)$$

Case 2: if $b \neq 0$ and $b^2 + 4ad - 4ac < 0$, then

$$\begin{aligned} \left(\frac{G'(\zeta)}{G(\zeta)} \right) &= \frac{b}{2(d-c)} \\ &+ \frac{\sqrt{4ac - 4ad - b^2}}{2(d-c)} \left(\frac{C_2 \cos \left(\frac{\zeta \sqrt{4ac - 4ad - b^2}}{2d} \right) - C_1 \sin \left(\frac{\zeta \sqrt{4ac - 4ad - b^2}}{2d} \right)}{C_1 \cos \left(\frac{\zeta \sqrt{4ac - 4ad - b^2}}{2d} \right) + C_2 \sin \left(\frac{\zeta \sqrt{4ac - 4ad - b^2}}{2d} \right)} \right). \end{aligned} \quad (31)$$

Case 3: if $b \neq 0$ and $b^2 + 4ad - 4ac = 0$, then

$$\left(\frac{G'(\zeta)}{G(\zeta)} \right) = \frac{b}{2(d-c)} + \frac{dD}{(d-c)(C-D\zeta)}. \quad (32)$$

Case 4: if $b = 0$ and $ad - ac > 0$, then

$$\left(\frac{G'(\zeta)}{G(\zeta)} \right) = \frac{\sqrt{ad - ac}}{(d-c)} \left(\frac{C_1 \sinh \left(\frac{\zeta \sqrt{ad - ac}}{d} \right) + C_2 \cosh \left(\frac{\zeta \sqrt{ad - ac}}{d} \right)}{C_1 \cosh \left(\frac{\zeta \sqrt{ad - ac}}{d} \right) + C_2 \sinh \left(\frac{\zeta \sqrt{ad - ac}}{d} \right)} \right). \quad (33)$$

Case 5: if $b = 0$ and $ad - ac < 0$, then

$$\left(\frac{G'(\zeta)}{G(\zeta)} \right) = \frac{\sqrt{ac - ad}}{d-c} \left(\frac{C_2 \cos \left(\frac{\zeta \sqrt{ac - ad}}{d} \right) - C_1 \sin \left(\frac{\zeta \sqrt{ac - ad}}{d} \right)}{C_1 \cos \left(\frac{\zeta \sqrt{ac - ad}}{d} \right) + C_2 \sin \left(\frac{\zeta \sqrt{ac - ad}}{d} \right)} \right). \quad (34)$$

where a, b, c, d, C_1 and C_2 are constants.

Step 5:

Substituting Eq. (28) along Eq. (29) into Eq. (27) and collecting coefficients of each power of $\left(\frac{G'(\zeta)}{G(\zeta)} \right)$. Putting each coefficient equal to zero, we achieve a set of algebraic equations involving $\nu, \kappa, \alpha_i, (i = 0, \pm 1, \pm 2, \dots, \pm m)$ and other parameters.

Step 6:

Solving the obtained set of equations by using Mathematica software.

Step 7:

Putting the gained solutions into Eq. (28) and we get exact solutions of Eq. (25).

4 Mathematical Treatment of the Model:

Let's suppose the travelling wave transform given as follows;

$$q(s, t) = Q(\zeta) \times \exp\left(\iota\left(\mu \frac{\Gamma(\varrho+1)}{\epsilon} s^\epsilon + \rho \frac{\Gamma(\varrho+1)}{\epsilon} t^\epsilon\right)\right), \quad \zeta = \lambda \frac{\Gamma(\varrho+1)}{\epsilon} s^\epsilon + \tau \frac{\Gamma(\varrho+1)}{\epsilon} t^\epsilon. \quad (35)$$

where $Q(\zeta)$ shows the amplitude of wave function while ρ and τ represent the time velocity. Parameters μ and λ are obtaining from asset price of the product.

Inserting Eq.(35) into Eq.(8), result in the form of real and imaginary parts given as follows:
Real part:

$$2\Omega Q^3 + \delta\lambda^2 Q'' - (\delta\mu^2 + 2\rho) Q = 0. \quad (36)$$

Imaginary part:

$$(\delta\mu\lambda + \tau)Q' = 0. \quad (37)$$

From Eq.(37), we get the velocity of wave function given as follows:

$$\tau = -\delta\mu\lambda. \quad (38)$$

Applying the homogenous balance method into Eq.(36), we get $m = 1$

Now we will gain the exact solutions of Eq.(36) by using above mentioned three methods.

5 Exact Solutions Through exp_a function Method:

For $m = 1$, Eq.(12) changes into:

$$Q(\zeta) = \frac{\alpha_0 + \alpha_1 d^\zeta}{\beta_0 + \beta_1 d^\zeta}. \quad (39)$$

Inserting Eq.(39) into Eq.(36) and solving the system of equations, we obtain different solution sets given as follows:

Set 1:

$$\left\{ \alpha_0 = -\frac{i\beta_0\sqrt{\delta}\lambda\log(d)}{2\sqrt{\Omega}}, \alpha_1 = \frac{i\beta_1\sqrt{\delta}\lambda\log(d)}{2\sqrt{\Omega}}, \rho = -\frac{1}{4}\delta(\lambda^2\log^2(d) + 2\mu^2) \right\}. \quad (40)$$

From Eqs.(40), (39) and (35), we get

$$q(s, t) = -\frac{i\sqrt{\delta}\lambda\log(d)}{2\sqrt{\Omega}} \left(\frac{\beta_0 - \beta_1 d^\zeta}{\beta_0 + \beta_1 d^\zeta} \right) \times \exp\left(\iota\left(\mu \frac{\Gamma(\varrho+1)}{\epsilon} s^\epsilon - \frac{1}{4}\delta(\lambda^2\log^2(d) + 2\mu^2) \frac{\Gamma(\varrho+1)}{\epsilon} t^\epsilon\right)\right). \quad (41)$$

Set 2:

$$\left\{ \alpha_0 = \frac{i\beta_0\sqrt{\delta}\lambda\log(d)}{2\sqrt{\Omega}}, \alpha_1 = -\frac{i\beta_1\sqrt{\delta}\lambda\log(d)}{2\sqrt{\Omega}}, \rho = -\frac{1}{4}\delta(\lambda^2\log^2(d) + 2\mu^2) \right\}. \quad (42)$$

From Eqs.(42), (39) and (35), we get

$$q(s, t) = \frac{i\sqrt{\delta}\lambda\log(d)}{2\sqrt{\Omega}} \left(\frac{\beta_0 - \beta_1 d^\zeta}{\beta_0 + \beta_1 d^\zeta} \right) \times \exp\left(\iota\left(\mu\frac{\Gamma(\varrho+1)}{\epsilon}s^\epsilon - \frac{1}{4}\delta(\lambda^2\log^2(d) + 2\mu^2)\frac{\Gamma(\varrho+1)}{\epsilon}t^\epsilon\right)\right). \quad (43)$$

where $\zeta = \lambda\frac{\Gamma(\varrho+1)}{\epsilon}s^\epsilon - \delta\mu\lambda\frac{\Gamma(\varrho+1)}{\epsilon}t^\epsilon$.

5.1 Exact Solutions Through EShGEEM:

For $m = 1$, Eq. (23), Eq. (24) and Eq. (16) become:

$$Q(\zeta) = \alpha_0 \pm \beta_1 \operatorname{csch}(\zeta) \pm \alpha_1 \operatorname{coth}(\zeta). \quad (44)$$

$$Q(\zeta) = \alpha_0 \pm \iota\beta_1 \operatorname{sech}(\zeta) \pm \alpha_1 \tanh(\zeta). \quad (45)$$

$$Q(\zeta) = \alpha_0 + \beta_1 \sinh(p) + \alpha_1 \cosh(p). \quad (46)$$

where α_0, α_1 and β_1 are unknowns. Substituting Eq. (46) into Eq. (36), we obtain the algebraic equations having $\alpha_0, \alpha_1, \beta_1$ and other parameters. Now with the help of software, we get different solution sets given as:

Set 1:

$$\left\{ \alpha_0 = 0, \alpha_1 = -\frac{i\sqrt{\delta}\lambda}{\sqrt{\Omega}}, \beta_1 = 0, \rho = -\frac{1}{2}\delta(2\lambda^2 + \mu^2) \right\}. \quad (47)$$

From Eqs.(47), (44) and (35), we get

$$q_1(s, t) = \mp \frac{i\sqrt{\delta}\lambda}{\sqrt{\Omega}} \operatorname{coth}(\zeta) \times \exp\left(\iota\left(\mu\frac{\Gamma(\varrho+1)}{\epsilon}s^\epsilon - \frac{1}{2}\delta(2\lambda^2 + \mu^2)\frac{\Gamma(\varrho+1)}{\epsilon}t^\epsilon\right)\right). \quad (48)$$

From Eqs.(47), (45) and (35), we get

$$q_2(s, t) = \mp \frac{i\sqrt{\delta}\lambda}{\sqrt{\Omega}} \tanh(\zeta) \times \exp\left(\iota\left(\mu\frac{\Gamma(\varrho+1)}{\epsilon}s^\epsilon - \frac{1}{2}\delta(2\lambda^2 + \mu^2)\frac{\Gamma(\varrho+1)}{\epsilon}t^\epsilon\right)\right). \quad (49)$$

Set 2:

$$\left\{ \alpha_0 = 0, \alpha_1 = \frac{i\sqrt{\delta}\lambda}{\sqrt{\Omega}}, \beta_1 = 0, \rho = -\frac{1}{2}\delta(2\lambda^2 + \mu^2) \right\}. \quad (50)$$

From Eqs.(50), (44) and (35), we get

$$q_1(s, t) = \pm \frac{i\sqrt{\delta}\lambda}{\sqrt{\Omega}} \coth(\zeta) \times \exp(\iota(\mu \frac{\Gamma(\varrho+1)}{\epsilon} s^\epsilon - \frac{1}{2}\delta (2\lambda^2 + \mu^2) \frac{\Gamma(\varrho+1)}{\epsilon} t^\epsilon)). \quad (51)$$

From Eqs.(50), (45) and (35), we get

$$q_2(s, t) = \pm \frac{i\sqrt{\delta}\lambda}{\sqrt{\Omega}} \tanh(\zeta) \times \exp(\iota(\mu \frac{\Gamma(\varrho+1)}{\epsilon} s^\epsilon - \frac{1}{2}\delta (2\lambda^2 + \mu^2) \frac{\Gamma(\varrho+1)}{\epsilon} t^\epsilon)). \quad (52)$$

Set 3:

$$\left\{ \alpha_0 = 0, \alpha_1 = -\frac{i\sqrt{\delta}\lambda}{2\sqrt{\Omega}}, \beta_1 = -\frac{i\sqrt{\delta}\lambda}{2\sqrt{\Omega}}, \rho = -\frac{1}{4}\delta (\lambda^2 + 2\mu^2) \right\}. \quad (53)$$

From Eqs.(53), (44) and (35), we get

$$q_1(s, t) = \mp \frac{i\sqrt{\delta}\lambda}{2\sqrt{\Omega}} (\coth(\zeta) + \operatorname{csch}(\zeta)) \times \exp(\iota(\mu \frac{\Gamma(\varrho+1)}{\epsilon} s^\epsilon - \frac{1}{4}\delta (\lambda^2 + 2\mu^2) \frac{\Gamma(\varrho+1)}{\epsilon} t^\epsilon)). \quad (54)$$

From Eqs.(53), (45) and (35), we get

$$q_2(s, t) = \mp \frac{i\sqrt{\delta}\lambda}{2\sqrt{\Omega}} (\iota \operatorname{sech}(\zeta) + \tanh(\zeta)) \times \exp(\iota(\mu \frac{\Gamma(\varrho+1)}{\epsilon} s^\epsilon - \frac{1}{4}\delta (\lambda^2 + 2\mu^2) \frac{\Gamma(\varrho+1)}{\epsilon} t^\epsilon)). \quad (55)$$

Set 4:

$$\left\{ \alpha_0 = 0, \alpha_1 = \frac{i\sqrt{\delta}\lambda}{2\sqrt{\Omega}}, \beta_1 = -\frac{i\sqrt{\delta}\lambda}{2\sqrt{\Omega}}, \rho = -\frac{1}{4}\delta (\lambda^2 + 2\mu^2) \right\}. \quad (56)$$

From Eqs.(56), (44) and (35), we get

$$q_1(s, t) = \frac{i\sqrt{\delta}\lambda}{2\sqrt{\Omega}} (\pm \coth(\zeta) \mp \operatorname{csch}(\zeta)) \times \exp(\iota(\mu \frac{\Gamma(\varrho+1)}{\epsilon} s^\epsilon - \frac{1}{4}\delta (\lambda^2 + 2\mu^2) \frac{\Gamma(\varrho+1)}{\epsilon} t^\epsilon)). \quad (57)$$

From Eqs.(56), (45) and (35), we get

$$q_2(s, t) = \frac{i\sqrt{\delta}\lambda}{2\sqrt{\Omega}} (\pm \tanh(\zeta) \mp \iota \operatorname{sech}(\zeta)) \times \exp(\iota(\mu \frac{\Gamma(\varrho+1)}{\epsilon} s^\epsilon - \frac{1}{4}\delta (\lambda^2 + 2\mu^2) \frac{\Gamma(\varrho+1)}{\epsilon} t^\epsilon)). \quad (58)$$

Set 5:

$$\left\{ \alpha_0 = 0, \alpha_1 = -\frac{i\sqrt{\delta}\lambda}{2\sqrt{\Omega}}, \beta_1 = \frac{i\sqrt{\delta}\lambda}{2\sqrt{\Omega}}, \rho = -\frac{1}{4}\delta (\lambda^2 + 2\mu^2) \right\}. \quad (59)$$

From Eqs.(59), (44) and (35), we get

$$q_1(s, t) = -\frac{i\sqrt{\delta}\lambda}{2\sqrt{\Omega}}(\pm \coth(\zeta) \mp \operatorname{csch}(\zeta)) \times \exp(\iota(\mu \frac{\Gamma(\varrho+1)}{\epsilon} s^\epsilon - \frac{1}{4}\delta(\lambda^2 + 2\mu^2) \frac{\Gamma(\varrho+1)}{\epsilon} t^\epsilon)). \quad (60)$$

From Eqs.(59), (45) and (35), we get

$$q_2(s, t) = -\frac{i\sqrt{\delta}\lambda}{2\sqrt{\Omega}}(\pm \tanh(\zeta) \mp \iota \operatorname{sech}(\zeta)) \times \exp(\iota(\mu \frac{\Gamma(\varrho+1)}{\epsilon} s^\epsilon - \frac{1}{4}\delta(\lambda^2 + 2\mu^2) \frac{\Gamma(\varrho+1)}{\epsilon} t^\epsilon)). \quad (61)$$

Set 6:

$$\left\{ \alpha_0 = 0, \alpha_1 = \frac{i\sqrt{\delta}\lambda}{2\sqrt{\Omega}}, \beta_1 = \frac{i\sqrt{\delta}\lambda}{2\sqrt{\Omega}}, \rho = -\frac{1}{4}\delta(\lambda^2 + 2\mu^2) \right\}. \quad (62)$$

From Eqs.(62), (44) and (35), we get

$$q_1(s, t) = \pm \frac{i\sqrt{\delta}\lambda}{2\sqrt{\Omega}}(\coth(\zeta) + \operatorname{csch}(\zeta)) \times \exp(\iota(\mu \frac{\Gamma(\varrho+1)}{\epsilon} s^\epsilon - \frac{1}{4}\delta(\lambda^2 + 2\mu^2) \frac{\Gamma(\varrho+1)}{\epsilon} t^\epsilon)). \quad (63)$$

From Eqs.(62), (45) and (35), we get

$$q_2(s, t) = \pm \frac{i\sqrt{\delta}\lambda}{2\sqrt{\Omega}}(\iota \operatorname{sech}(\zeta) + \tanh(\zeta)) \times \exp(\iota(\mu \frac{\Gamma(\varrho+1)}{\epsilon} s^\epsilon - \frac{1}{4}\delta(\lambda^2 + 2\mu^2) \frac{\Gamma(\varrho+1)}{\epsilon} t^\epsilon)). \quad (64)$$

Set 7:

$$\left\{ \alpha_0 = 0, \alpha_1 = 0, \beta_1 = -\frac{i\sqrt{\delta}\lambda}{\sqrt{\Omega}}, \rho = \frac{1}{2}\delta(\lambda^2 - \mu^2) \right\}. \quad (65)$$

From Eqs.(65), (44) and (35), we get

$$q_1(s, t) = \mp \frac{i\sqrt{\delta}\lambda}{\sqrt{\Omega}}\operatorname{csch}(\zeta) \times \exp(\iota(\mu \frac{\Gamma(\varrho+1)}{\epsilon} s^\epsilon + \frac{1}{2}\delta(\lambda^2 - \mu^2) \frac{\Gamma(\varrho+1)}{\epsilon} t^\epsilon)). \quad (66)$$

From Eqs.(65), (45) and (35), we get

$$q_2(s, t) = \pm \frac{\sqrt{\delta}\lambda}{\sqrt{\Omega}}\operatorname{sech}(\zeta) \times \exp(\iota(\mu \frac{\Gamma(\varrho+1)}{\epsilon} s^\epsilon + \frac{1}{2}\delta(\lambda^2 - \mu^2) \frac{\Gamma(\varrho+1)}{\epsilon} t^\epsilon)). \quad (67)$$

Set 8:

$$\left\{ \alpha_0 = 0, \alpha_1 = 0, \beta_1 = \frac{i\sqrt{\delta}\lambda}{\sqrt{\Omega}}, \rho = \frac{1}{2}\delta(\lambda^2 - \mu^2) \right\}. \quad (68)$$

From Eqs.(68), (44) and (35), we get

$$q_1(s, t) = \pm \frac{i\sqrt{\delta}\lambda}{\sqrt{\Omega}}\operatorname{csch}(\zeta) \times \exp(\iota(\mu \frac{\Gamma(\varrho+1)}{\epsilon} s^\epsilon + \frac{1}{2}\delta(\lambda^2 - \mu^2) \frac{\Gamma(\varrho+1)}{\epsilon} t^\epsilon)). \quad (69)$$

From Eqs.(68), (45) and (35), we get

$$q_2(s, t) = \mp \frac{\sqrt{\delta}\lambda}{\sqrt{\Omega}} \text{sech}(\zeta) \times \exp(\iota(\mu \frac{\Gamma(\varrho+1)}{\epsilon} s^\epsilon + \frac{1}{2} \delta (\lambda^2 - \mu^2) \frac{\Gamma(\varrho+1)}{\epsilon} t^\epsilon)), \quad (70)$$

where $\zeta = \lambda \frac{\Gamma(\varrho+1)}{\epsilon} s^\epsilon - \delta \mu \lambda \frac{\Gamma(\varrho+1)}{\epsilon} t^\epsilon$.

5.2 Exact Solutions Through Extended (G'/G) -expansion Method:

For $m = 1$, Eq. (28) becomes:

$$Q(\zeta) = \alpha_{-1} \left(\frac{G'(\zeta)}{G(\zeta)} \right)^{-1} + \alpha_0 + \alpha_1 \left(\frac{G'(\zeta)}{G(\zeta)} \right), \quad (71)$$

where α_{-1}, α_0 and α_1 are unknowns.

Inserting Eq. (71) along Eq. (29) into Eq. (36) and solving the system for $\alpha_{-1}, \alpha_0, \alpha_1$ and other parameters, we gain different solution sets given as:

Set 1:

$$\left\{ \alpha_{-1} = -\frac{ia\sqrt{\delta}\lambda}{d\sqrt{\Omega}}, \alpha_0 = -\frac{ib\sqrt{\delta}\lambda}{2d\sqrt{\Omega}}, \alpha_1 = 0, \rho = -\frac{\delta(4a\lambda^2(d-c) + b^2\lambda^2 + 2d^2\mu^2)}{4d^2} \right\}. \quad (72)$$

From Eqs.(72), (71), (30) and (35), we get

$$\begin{aligned} q(s, t) = & -\frac{i\sqrt{\delta}\lambda}{d\sqrt{\Omega}} \left(\frac{b}{2} + a \left(\frac{b}{2(d-c)} \right. \right. \\ & + \frac{\sqrt{-4ac+4ad+b^2}}{2(d-c)} \left(\frac{C_1 \sinh(\frac{\zeta\sqrt{-4ac+4ad+b^2}}{2d}) + C_2 \cosh(\frac{\zeta\sqrt{-4ac+4ad+b^2}}{2d})}{C_1 \cosh(\frac{\zeta\sqrt{-4ac+4ad+b^2}}{2d}) + C_2 \sinh(\frac{\zeta\sqrt{-4ac+4ad+b^2}}{2d})} \right)^{-1} \\ & \times \exp(\iota(\mu \frac{\Gamma(\varrho+1)}{\epsilon} s^\epsilon - \frac{\delta(4a\lambda^2(d-c) + b^2\lambda^2 + 2d^2\mu^2)}{4d^2} \frac{\Gamma(\varrho+1)}{\epsilon} t^\epsilon)). \end{aligned} \quad (73)$$

From Eqs.(72), (71), (31) and (35), we get

$$\begin{aligned} q(s, t) = & -\frac{i\sqrt{\delta}\lambda}{d\sqrt{\Omega}} \left(\frac{b}{2} + a \left(\frac{b}{2(d-c)} \right. \right. \\ & + \frac{\sqrt{4ac-4ad-b^2}}{2(d-c)} \left(\frac{C_2 \cos\left(\frac{\zeta\sqrt{4ac-4ad-b^2}}{2d}\right) - C_1 \sin\left(\frac{\zeta\sqrt{4ac-4ad-b^2}}{2d}\right)}{C_1 \cos\left(\frac{\zeta\sqrt{4ac-4ad-b^2}}{2d}\right) + C_2 \sin\left(\frac{\zeta\sqrt{4ac-4ad-b^2}}{2d}\right)} \right)^{-1} \\ & \times \exp(\iota(\mu \frac{\Gamma(\varrho+1)}{\epsilon} s^\epsilon - \frac{\delta(4a\lambda^2(d-c) + b^2\lambda^2 + 2d^2\mu^2)}{4d^2} \frac{\Gamma(\varrho+1)}{\epsilon} t^\epsilon)). \end{aligned} \quad (74)$$

From Eqs.(72), (71), (33) and (35), we get

$$q(s, t) = -\frac{ia\sqrt{\delta}\lambda}{d\sqrt{\Omega}} \left(\frac{\sqrt{ad-ac}}{(d-c)} \left(\frac{C_1 \sinh\left(\frac{\zeta\sqrt{ad-ac}}{d}\right) + C_2 \cosh\left(\frac{\zeta\sqrt{ad-ac}}{d}\right)}{C_1 \cosh\left(\frac{\zeta\sqrt{ad-ac}}{d}\right) + C_2 \sinh\left(\frac{\zeta\sqrt{ad-ac}}{d}\right)} \right) \right)^{-1} \\ \times \exp\left(\iota\left(\mu\frac{\Gamma(\varrho+1)}{\epsilon} s^\epsilon - \frac{\delta(4a\lambda^2(d-c) + 2d^2\mu^2)}{4d^2} \frac{\Gamma(\varrho+1)}{\epsilon} t^\epsilon\right)\right). \quad (75)$$

From Eqs.(72), (71), (34) and (35), we get

$$q(s, t) = -\frac{ia\sqrt{\delta}\lambda}{d\sqrt{\Omega}} \left(\frac{\sqrt{ac-ad}}{d-c} \left(\frac{C_2 \cos\left(\frac{\zeta\sqrt{ac-ad}}{d}\right) - C_1 \sin\left(\frac{\zeta\sqrt{ac-ad}}{d}\right)}{C_1 \cos\left(\frac{\zeta\sqrt{ac-ad}}{d}\right) + C_2 \sin\left(\frac{\zeta\sqrt{ac-ad}}{d}\right)} \right) \right)^{-1} \\ \times \exp\left(\iota\left(\mu\frac{\Gamma(\varrho+1)}{\epsilon} s^\epsilon - \frac{\delta(4a\lambda^2(d-c) + 2d^2\mu^2)}{4d^2} \frac{\Gamma(\varrho+1)}{\epsilon} t^\epsilon\right)\right). \quad (76)$$

Set 2:

$$\left\{ \alpha_{-1} = \frac{ia\sqrt{\delta}\lambda}{d\sqrt{\Omega}}, \alpha_0 = \frac{ib\sqrt{\delta}\lambda}{2d\sqrt{\Omega}}, \alpha_1 = 0, \rho = -\frac{\delta(4a\lambda^2(d-c) + b^2\lambda^2 + 2d^2\mu^2)}{4d^2} \right\}. \quad (77)$$

From Eqs.(77), (71), (30) and (35), we get

$$q(s, t) = \frac{i\sqrt{\delta}\lambda}{d\sqrt{\Omega}} \left(\frac{b}{2} + a\left(\frac{b}{2(d-c)} \right. \right. \\ \left. \left. + \frac{\sqrt{-4ac+4ad+b^2}}{2(d-c)} \left(\frac{C_1 \sinh\left(\frac{\zeta\sqrt{-4ac+4ad+b^2}}{2d}\right) + C_2 \cosh\left(\frac{\zeta\sqrt{-4ac+4ad+b^2}}{2d}\right)}{C_1 \cosh\left(\frac{\zeta\sqrt{-4ac+4ad+b^2}}{2d}\right) + C_2 \sinh\left(\frac{\zeta\sqrt{-4ac+4ad+b^2}}{2d}\right)} \right) \right)^{-1} \right) \\ \times \exp\left(\iota\left(\mu\frac{\Gamma(\varrho+1)}{\epsilon} s^\epsilon - \frac{\delta(4a\lambda^2(d-c) + b^2\lambda^2 + 2d^2\mu^2)}{4d^2} \frac{\Gamma(\varrho+1)}{\epsilon} t^\epsilon\right)\right). \quad (78)$$

From Eqs.(77), (71), (31) and (35), we get

$$q(s, t) = \frac{i\sqrt{\delta}\lambda}{d\sqrt{\Omega}} \left(\frac{b}{2} + a\left(\frac{b}{2(d-c)} \right. \right. \\ \left. \left. + \frac{\sqrt{4ac-4ad-b^2}}{2(d-c)} \left(\frac{C_2 \cos\left(\frac{\zeta\sqrt{4ac-4ad-b^2}}{2d}\right) - C_1 \sin\left(\frac{\zeta\sqrt{4ac-4ad-b^2}}{2d}\right)}{C_1 \cos\left(\frac{\zeta\sqrt{4ac-4ad-b^2}}{2d}\right) + C_2 \sin\left(\frac{\zeta\sqrt{4ac-4ad-b^2}}{2d}\right)} \right) \right)^{-1} \right) \\ \times \exp\left(\iota\left(\mu\frac{\Gamma(\varrho+1)}{\epsilon} s^\epsilon - \frac{\delta(4a\lambda^2(d-c) + b^2\lambda^2 + 2d^2\mu^2)}{4d^2} \frac{\Gamma(\varrho+1)}{\epsilon} t^\epsilon\right)\right). \quad (79)$$

From Eqs.(77), (71), (33) and (35), we get

$$q(s, t) = \frac{ia\sqrt{\delta}\lambda}{d\sqrt{\Omega}} \left(\frac{\sqrt{ad-ac}}{(d-c)} \left(\frac{C_1 \sinh\left(\frac{\zeta\sqrt{ad-ac}}{d}\right) + C_2 \cosh\left(\frac{\zeta\sqrt{ad-ac}}{d}\right)}{C_1 \cosh\left(\frac{\zeta\sqrt{ad-ac}}{d}\right) + C_2 \sinh\left(\frac{\zeta\sqrt{ad-ac}}{d}\right)} \right) \right)^{-1} \\ \times \exp\left(\iota\left(\mu\frac{\Gamma(\varrho+1)}{\epsilon}s^\epsilon - \frac{\delta(4a\lambda^2(d-c) + 2d^2\mu^2)}{4d^2}\frac{\Gamma(\varrho+1)}{\epsilon}t^\epsilon\right)\right). \quad (80)$$

From Eqs.(77), (71), (34) and (35), we get

$$q(s, t) = \frac{ia\sqrt{\delta}\lambda}{d\sqrt{\Omega}} \left(\frac{\sqrt{ac-ad}}{d-c} \left(\frac{C_2 \cos\left(\frac{\zeta\sqrt{ac-ad}}{d}\right) - C_1 \sin\left(\frac{\zeta\sqrt{ac-ad}}{d}\right)}{C_1 \cos\left(\frac{\zeta\sqrt{ac-ad}}{d}\right) + C_2 \sin\left(\frac{\zeta\sqrt{ac-ad}}{d}\right)} \right) \right)^{-1} \\ \times \exp\left(\iota\left(\mu\frac{\Gamma(\varrho+1)}{\epsilon}s^\epsilon - \frac{\delta(4a\lambda^2(d-c) + 2d^2\mu^2)}{4d^2}\frac{\Gamma(\varrho+1)}{\epsilon}t^\epsilon\right)\right). \quad (81)$$

Set 3:

$$\left\{ \alpha_{-1} = 0, \alpha_0 = -\frac{ib\sqrt{\delta}\lambda}{2d\sqrt{\Omega}}, \alpha_1 = -\frac{i\sqrt{\delta}\lambda(c-d)}{d\sqrt{\Omega}}, \rho = -\frac{\delta(4a\lambda^2(d-c) + b^2\lambda^2 + 2d^2\mu^2)}{4d^2} \right\}. \quad (82)$$

From Eqs.(82), (71), (30) and (35), we get

$$q(s, t) = \frac{-i\sqrt{\delta}\lambda}{d\sqrt{\Omega}} \left(\frac{b}{2} - \left(\frac{b}{2} + \frac{\sqrt{-4ac+4ad+b^2}}{2} \left(\frac{C_1 \sinh\left(\frac{\zeta\sqrt{-4ac+4ad+b^2}}{2d}\right) + C_2 \cosh\left(\frac{\zeta\sqrt{-4ac+4ad+b^2}}{2d}\right)}{C_1 \cosh\left(\frac{\zeta\sqrt{-4ac+4ad+b^2}}{2d}\right) + C_2 \sinh\left(\frac{\zeta\sqrt{-4ac+4ad+b^2}}{2d}\right)} \right) \right) \right) \\ \times \exp\left(\iota\left(\mu\frac{\Gamma(\varrho+1)}{\epsilon}s^\epsilon - \frac{\delta(4a\lambda^2(d-c) + b^2\lambda^2 + 2d^2\mu^2)}{4d^2}\frac{\Gamma(\varrho+1)}{\epsilon}t^\epsilon\right)\right). \quad (83)$$

From Eqs.(82), (71), (31) and (35), we get

$$q(s, t) = -\frac{i\sqrt{\delta}\lambda}{d\sqrt{\Omega}} \left(\frac{b}{2} + (c-d) \left(\frac{b}{2(d-c)} + \frac{\sqrt{4ac-4ad-b^2}}{2(d-c)} \left(\frac{C_2 \cos\left(\frac{\zeta\sqrt{4ac-4ad-b^2}}{2d}\right) - C_1 \sin\left(\frac{\zeta\sqrt{4ac-4ad-b^2}}{2d}\right)}{C_1 \cos\left(\frac{\zeta\sqrt{4ac-4ad-b^2}}{2d}\right) + C_2 \sin\left(\frac{\zeta\sqrt{4ac-4ad-b^2}}{2d}\right)} \right) \right) \right) \\ \times \exp\left(\iota\left(\mu\frac{\Gamma(\varrho+1)}{\epsilon}s^\epsilon - \frac{\delta(4a\lambda^2(d-c) + b^2\lambda^2 + 2d^2\mu^2)}{4d^2}\frac{\Gamma(\varrho+1)}{\epsilon}t^\epsilon\right)\right). \quad (84)$$

From Eqs.(82), (71), (33) and (35), we get

$$q(s, t) = -\frac{i(c-d)\sqrt{\delta}\lambda}{d\sqrt{\Omega}} \left(\frac{\sqrt{ad-ac}}{(d-c)} \left(\frac{C_1 \sinh\left(\frac{\zeta\sqrt{ad-ac}}{d}\right) + C_2 \cosh\left(\frac{\zeta\sqrt{ad-ac}}{d}\right)}{C_1 \cosh\left(\frac{\zeta\sqrt{ad-ac}}{d}\right) + C_2 \sinh\left(\frac{\zeta\sqrt{ad-ac}}{d}\right)} \right) \right) \\ \times \exp\left(\iota\left(\mu\frac{\Gamma(\varrho+1)}{\epsilon}s^\epsilon - \frac{\delta(4a\lambda^2(d-c) + 2d^2\mu^2)}{4d^2}\frac{\Gamma(\varrho+1)}{\epsilon}t^\epsilon\right)\right). \quad (85)$$

From Eqs.(82), (71), (34) and (35), we get

$$q(s, t) = -\frac{i(c-d)\sqrt{\delta}\lambda}{d\sqrt{\Omega}} \left(\frac{\sqrt{ac-ad}}{d-c} \left(\frac{C_2 \cos\left(\frac{\zeta\sqrt{ac-ad}}{d}\right) - C_1 \sin\left(\frac{\zeta\sqrt{ac-ad}}{d}\right)}{C_1 \cos\left(\frac{\zeta\sqrt{ac-ad}}{d}\right) + C_2 \sin\left(\frac{\zeta\sqrt{ac-ad}}{d}\right)} \right) \right) \\ \times \exp\left(\iota\left(\mu\frac{\Gamma(\varrho+1)}{\epsilon}s^\epsilon - \frac{\delta(4a\lambda^2(d-c) + 2d^2\mu^2)}{4d^2}\frac{\Gamma(\varrho+1)}{\epsilon}t^\epsilon\right)\right). \quad (86)$$

Set 4:

$$\left\{ \alpha_{-1} = 0, \alpha_0 = \frac{ib\sqrt{\delta}\lambda}{2d\sqrt{\Omega}}, \alpha_1 = \frac{i\sqrt{\delta}\lambda(c-d)}{d\sqrt{\Omega}}, \rho = -\frac{\delta(4a\lambda^2(d-c) + b^2\lambda^2 + 2d^2\mu^2)}{4d^2} \right\}. \quad (87)$$

From Eqs.(87), (71), (30) and (35), we get

$$q(s, t) = \frac{i\sqrt{\delta}\lambda}{d\sqrt{\Omega}} \left(\frac{b}{2} + (c-d)\left(\frac{b}{2(d-c)} \right. \right. \\ \left. \left. + \frac{\sqrt{-4ac+4ad+b^2}}{2(d-c)} \left(\frac{C_1 \sinh\left(\frac{\zeta\sqrt{-4ac+4ad+b^2}}{2d}\right) + C_2 \cosh\left(\frac{\zeta\sqrt{-4ac+4ad+b^2}}{2d}\right)}{C_1 \cosh\left(\frac{\zeta\sqrt{-4ac+4ad+b^2}}{2d}\right) + C_2 \sinh\left(\frac{\zeta\sqrt{-4ac+4ad+b^2}}{2d}\right)} \right) \right) \right) \\ \times \exp\left(\iota\left(\mu\frac{\Gamma(\varrho+1)}{\epsilon}s^\epsilon - \frac{\delta(4a\lambda^2(d-c) + b^2\lambda^2 + 2d^2\mu^2)}{4d^2}\frac{\Gamma(\varrho+1)}{\epsilon}t^\epsilon\right)\right). \quad (88)$$

From Eqs.(87), (71), (31) and (35), we get

$$q(s, t) = \frac{i\sqrt{\delta}\lambda}{d\sqrt{\Omega}} \left(\frac{b}{2} + (c-d)\left(\frac{b}{2(d-c)} \right. \right. \\ \left. \left. + \frac{\sqrt{4ac-4ad-b^2}}{2(d-c)} \left(\frac{C_2 \cos\left(\frac{\zeta\sqrt{4ac-4ad-b^2}}{2d}\right) - C_1 \sin\left(\frac{\zeta\sqrt{4ac-4ad-b^2}}{2d}\right)}{C_1 \cos\left(\frac{\zeta\sqrt{4ac-4ad-b^2}}{2d}\right) + C_2 \sin\left(\frac{\zeta\sqrt{4ac-4ad-b^2}}{2d}\right)} \right) \right) \right) \\ \times \exp\left(\iota\left(\mu\frac{\Gamma(\varrho+1)}{\epsilon}s^\epsilon - \frac{\delta(4a\lambda^2(d-c) + b^2\lambda^2 + 2d^2\mu^2)}{4d^2}\frac{\Gamma(\varrho+1)}{\epsilon}t^\epsilon\right)\right). \quad (89)$$

From Eqs.(87), (71), (33) and (35), we get

$$q(s, t) = \frac{i(c-d)\sqrt{\delta}\lambda}{d\sqrt{\Omega}} \left(\frac{\sqrt{ad-ac}}{(d-c)} \left(\frac{C_1 \sinh\left(\frac{\zeta\sqrt{ad-ac}}{d}\right) + C_2 \cosh\left(\frac{\zeta\sqrt{ad-ac}}{d}\right)}{C_1 \cosh\left(\frac{\zeta\sqrt{ad-ac}}{d}\right) + C_2 \sinh\left(\frac{\zeta\sqrt{ad-ac}}{d}\right)} \right) \right) \\ \times \exp\left(\iota\left(\mu\frac{\Gamma(\varrho+1)}{\epsilon}s^\epsilon - \frac{\delta(4a\lambda^2(d-c) + 2d^2\mu^2)}{4d^2}\frac{\Gamma(\varrho+1)}{\epsilon}t^\epsilon\right)\right). \quad (90)$$

From Eqs.(87), (71), (34) and (35), we get

$$q(s, t) = \frac{i(c-d)\sqrt{\delta}\lambda}{d\sqrt{\Omega}} \left(\frac{\sqrt{ac-ad}}{d-c} \left(\frac{C_2 \cos\left(\frac{\zeta\sqrt{ac-ad}}{d}\right) - C_1 \sin\left(\frac{\zeta\sqrt{ac-ad}}{d}\right)}{C_1 \cos\left(\frac{\zeta\sqrt{ac-ad}}{d}\right) + C_2 \sin\left(\frac{\zeta\sqrt{ac-ad}}{d}\right)} \right) \right) \\ \times \exp\left(\iota\left(\mu\frac{\Gamma(\varrho+1)}{\epsilon}s^\epsilon - \frac{\delta(4a\lambda^2(d-c) + 2d^2\mu^2)}{4d^2}\frac{\Gamma(\varrho+1)}{\epsilon}t^\epsilon\right)\right). \quad (91)$$

where $\zeta = \lambda\frac{\Gamma(\varrho+1)}{\epsilon}s^\epsilon - \delta\mu\lambda\frac{\Gamma(\varrho+1)}{\epsilon}t^\epsilon$ for all above mentioned solutions.

6 Illustrations with graphics

Here, we show some two-dimensional and three-dimensional figures to help clarify the solutions we presented. Figures 1-3 depict some of the analytical solutions. In Figure 1, we use our method to introduce the graph of (41) at $\delta = 0.5, \lambda = 0.3, \varrho = 0.1, \mu = 2, \Omega = 0.7, \beta_0 = 0.1, \beta_1 = 0.1, d = 0.1, \epsilon = 1$. In addition, Figure 2 shows the graph of (48) $\delta = 0.5, \lambda = 0.1, \varrho = 0.5, \mu = 1, \Omega = 0.1, \epsilon = 1$. Finally, the graph of (73) at $\delta = 0.3, \lambda = 0.4, \varrho = 0.5, \mu = 6, \Omega = 0.4, d = 0.17, a = 0.1, c = 0.01, b = 0.4, C_1 = 0.4, C_2 = 0.5$ presented in Figure 3.

7 Conclusion

We have succeed to obtain the modernistic exact solutions of M-fractional Ivancevic option pricing model by utilizing exp_a function, extended Sinh-Gordon equation expansion and extended (G'/G) -Expansion methods. The gained solutions are also verified and demonstrated through graphs by using MATHEMATICA software. The obtained results are also explained graphically by 2-D,3-D and contour plots. Finally, it is suggested, to deal the other non-linear PDEs, the exp_a function, extended Sinh-Gordon equation expansion and extended (G'/G) -Expansion methods are very helpful, reliable and straight forward. Results achieved in this paper may useful for the progress in the supplementary analyzing of this model. Fractional derivatives (local, conformable, truncated M-fractional, beta fractional, caputo-Fabrizio fractional derivative,...) have attracted extensive attention in the field of mathematical physics. How to apply the proposed methods to study the fractional PDEs will be the focus of our future research.




Figure 1: Structure of (41) at $\delta = 0.5, \lambda = 0.3, \varrho = 0.1, \mu = 2, \Omega = 0.7, \beta_0 = 0.1, \beta_1 = 0.1, d = 0.1, \epsilon = 1$




Figure 2: Structure of (48) at $\delta = 0.5, \lambda = 0.1, \varrho = 0.5, \mu = 1, \Omega = 0.1, \epsilon = 1$




Figure 3: Structure of (73) at $\delta = 0.3, \lambda = 0.4, \varrho = 0.5, \mu = 6, \Omega = 0.4, d = 0.17, a = 0.1, c = 0.01, b = 0.4, C_1 = 0.4, C_2 = 0.5$.

References

- [1] Ghanbari, Behzad and Osman, MS and Baleanu, Dumitru; Generalized exponential rational function method for extended Zakharov–Kuzetsov equation with conformable derivative, *Modern Physics Letters A*, 34, 20, 1950155, (2019).
- [2] Osman, MS and Ghanbari, Behzad; New optical solitary wave solutions of Fokas–Lenells equation in presence of perturbation terms by a novel approach, *Optik*, 175, 328–333, (2018).
- [3] Neirameh, Ahmad and Eslami, Mostafa; New solitary wave solutions for fractional Jaulent–Miodek hierarchy equation, *Modern Physics Letters B*, 2150612, (2022).
- [4] Rezazadeh, Hadi and Osman, Mohamed S and Eslami, Mostafa and Ekici, Mehmet and Sonmezoglu, Abdullah and Asma, Mir and Othman, Wan Ainun Mior and Wong, BR and Mirzazadeh, Mohammad and Zhou, Qin and others; Mitigating Internet bottleneck with fractional temporal evolution of optical solitons having quadratic–cubic nonlinearity, *Optik*, 164, 84–92, (2018).
- [5] Osman, MS; One-soliton shaping and inelastic collision between double solitons in the fifth-order variable-coefficient Sawada–Kotera equation, *Nonlinear Dynamics*, 96, 2, 1491–1496, (2019).
- [6] Ali, Khaled K and Wazwaz, Abdul-Majid and Osman, MS; Optical soliton solutions to the generalized nonautonomous nonlinear Schrödinger equations in optical fibers via the sine-Gordon expansion method, *Optik*, 208, 164132, (2020).

- [7] Aktar, Most Shewly and Akbar, M Ali and Osman, MS; Spatio-temporal dynamic solitary wave solutions and diffusion effects to the nonlinear diffusive predator-prey system and the diffusion-reaction equations, *Chaos, Solitons & Fractals*, 160, 112212, (2022).
- [8] Inan, Bilge and Osman, Mohamed S and Ak, Turgut and Baleanu, Dumitru; Analytical and numerical solutions of mathematical biology models: The Newell-Whitehead-Segel and Allen-Cahn equations, *Mathematical methods in the applied sciences*, 43, 5, 2588–2600, (2020).
- [9] Adel, M and Baleanu, Dumitru and Sadiya, Umme and Arefin, Mohammad Asif and Uddin, M Hafiz and Elamin, Mahjoub A and Osman, MS; Inelastic soliton wave solutions with different geometrical structures to fractional order nonlinear evolution equations, *Results in Physics*, 105661, (2022)
- [10] Mohamed, R.Ali and Ma, Wen-Xiu and Sadat, R; Lie symmetry analysis and invariant solutions for (2+1) dimensional Bogoyavlensky-Konopelchenko equation with variable-coefficient in wave propagation, *Journal of Ocean Engineering and Science*, 7(3), 248-254, (2022).
- [11] Ali, Karmina K and Yilmazer, Resat and Osman, MS; Dynamic behavior of the (3+1)-dimensional KdV–Calogero–Bogoyavlenskii–Schiff equation, *Optical and Quantum Electronics*, 54, 3, 1–15, (2022).
- [12] Tarla, Sibel and Ali, Karmina K and Yilmazer, Resat and Osman, MS; The dynamic behaviors of the Radhakrishnan–Kundu–Lakshmanan equation by Jacobi elliptic function expansion technique, *Optical and Quantum Electronics*, 54, 5, 1–12, (2022).
- [13] Rashid, Saima and Kubra, Khadija Tul and Sultana, Sobia and Agarwal, Praveen and Osman, MS; An approximate analytical view of physical and biological models in the setting of Caputo operator via Elzaki transform decomposition method, *Journal of Computational and Applied Mathematics*, 114378, (2022).
- [14] Ismael, Hajar F and Okumuş, İsmail and Aktürk, Tolga and Bulut, Hasan and Osman, MS; Analyzing study for the 3D potential Yu–Toda–Sasa–Fukuyama equation in the two-layer liquid medium, *Journal of Ocean Engineering and Science*, (2022).
- [15] Muhannad, A.Shallal and Ali, Khalid K and Raslan, Kamal and Rezazadeh, Hadi and Bekir, Ahmet; Exact solutions of the conformable fractional EW and MEW equations by a new generalized expansion method, *Journal of Ocean Engineering and Science*, 5(3), 223-229, (2020).
- [16] El-Ganaini, Shoukry and Al-Amr, Mohammed O; New abundant solitary wave structures for a variety of some nonlinear models of surface wave propagation with their geometric interpretations, *Mathematical Methods in the Applied Sciences*, (2022).

- [17] Al-Amr, Mohammed O and Rezazadeh, Hadi and Ali, Khalid K and Korkmaz, Alper; N1-soliton solution for Schrödinger equation with competing weakly nonlocal and parabolic law nonlinearities, Communications in Theoretical Physics, 72, 6, 065503, (2020).
- [18] Rasheed, Noha M and Al-Amr, Mohammed O and Az-Zo'bi, Emad A and Tash-toush, Mohammad A and Akinyemi, Lanre; Stable optical solitons for the Higher-order Non-Kerr NLSE via the modified simple equation method, Mathematics, 9, 16, 1986, (2021).
- [19] Eslami, Mostafa and Rezazadeh, Hadi; The first integral method for Wu–Zhang system with conformable time-fractional derivative, Calcolo, 53, 3, 475–485, (2016).
- [20] Rezazadeh, Hadi and Kumar, Dipankar and Neirameh, Ahmad and Eslami, Mostafa and Mirzazadeh, Mohammad; Applications of three methods for obtaining optical soliton solutions for the Lakshmanan–Porsezian–Daniel model with Kerr law nonlinearity, Pramana, 94, 1, 1–11, (2020).
- [21] Zafar, A and Raheel, M and Mirzazadeh, M and Eslami, M; Different soliton solutions to the modified equal-width wave equation with Beta-time fractional derivative via two different methods, Revista Mexicana de Fisica, 68, 1, (2022).
- [22] Tukur Abdulkadir Sulaiman, Gulnur Yel and Hasan Bulut; M-fractional solitons and periodic wave solutions to the Hirota–Maccari system, Modern Physics Letters B, 1950052, (2019).
- [23] J. Vanterler D A C. Sousa, and E. Capelas D E Oliveira; A new truncated M-fractional derivative type unifying some fractional derivative types with classical properties, International Journal of Analysis and Applications, 16, 1 , 83-96, (2018).
- [24] Ivancevic, V.G.: Adaptive-Wave Alternative for the Black-Scholes Option Pricing Model. Cogn. Comput. 2, 17–30,(2010).
- [25] A. T. Ali and E.R. Hassan, General Exp_a -function method for nonlinear evolution equations, Applied Mathematics and Computation, 217(2) 451-459 (2010).
- [26] E.M.E. Zayed and A.G. Al-Nowehy, Generalized kudryashov method and general exp_a function method for solving a high order nonlinear schrödinger equation, J. Space Explor, 6, 1-26 (2017).
- [27] K. Hosseini, Z. Ayati and R. Ansari, New exact solutions of the Tzitzéica-type equations in non-linear optics using the exp_a function method, Journal of Modern Optics, 65(7), 847-851 (2018).

- [28] A. Zafar, The \exp_a function method and the conformable time-fractional KdV equations, Nonlinear Engineering, 8, 728-732 (2019).
- [29] X. L. Yang, J. S. Tang; Travelling wave solutions for Konopelchenko-Dubrovsky equation using an extended sinh-Gordon equation expansion method; Communications in Theoretical Physics; 50; 10471051; (2008).
- [30] S. Sahoo, S. Saha Ray, M. A. Abdou, New exact solutions for time-fractional Kaup-Kupershmidt equation using improved (G'/G) -expansion and extended (G'/G) -expansion methods, Alexandria Engineering Journal, <https://doi.org/10.1016/j.aej.2020.06.043>, (2020).
- [31] Zafar, Asim and Ali, Khalid K and Raheel, Muhammad and Nisar, Kottakkaran Sooppy and Bekir, Ahmet; Abundant M-fractional optical solitons to the perturbed Gerdjikov-Ivanov equation treating the mathematical nonlinear optics, Optical and Quantum Electronics, 54, 1, 1–17, (2022).
- [32] Zafar, Asim and Bekir, Ahmet and Raheel, Muhammad and Rezazadeh, Hadi; Investigation for optical soliton solutions of two nonlinear Schrödinger equations via two concrete finite series methods, International Journal of Applied and Computational Mathematics, 6, 3, 1–13, (2020).
- [33] Seadawy, Aly R and Kumar, Dipankar and Chakrabarty, Anuz Kumar; Dispersive optical soliton solutions for the hyperbolic and cubic-quintic nonlinear Schrödinger equations via the extended sinh-Gordon equation expansion method, The European Physical Journal Plus, 133, 5, 182, (2018).
- [34] Bezgabadi, A Safaei and Bolorizadeh, MA; Analytic combined bright-dark, bright and dark solitons solutions of generalized nonlinear Schrödinger equation using extended sinh-Gordon equation expansion method, Results in Physics, 30, 104852, (2021).
- [35] Taghizadeh, Nasir and Moosavi Noori, Seyyedeh R and Moosavi Noori, Seyyedeh B; Application of the Extended (G'/G) -expansion Method to the Improved Eckhaus Equation, Applications and Applied Mathematics: An International Journal (AAM), 9, 1, 24, (2014).
- [36] Ekici, Mehmet; Soliton and other solutions of nonlinear time fractional parabolic equations using extended (G'/G) -expansion method, Optik, 130, 1312–1319, (2017).
- [37] Wu, Wenqing and Ma, Xin and Zeng, Bo and Zhang, Hui and Zhang, Peng; A novel multivariate grey system model with conformable fractional derivative and its applications, Computers & Industrial Engineering, 164, 107888, (2022).
- [38] R. Khalil and M. Al Horani and A. Yousef and M. Sababheh, A new defnition of fractional derivative, J.Comput.Appl.Math., 264, 65-70, (2014).

- [39] Nisar, Kottakkaran Sooppy and Ciancio, Armando and Ali, Khalid K and Osman, MS and Cattani, Carlo and Baleanu, Dumitru and Zafar, Asim and Raheel, M and Azeem, M; On beta-time fractional biological population model with abundant solitary wave structures, *Alexandria Engineering Journal*, 61, 3, 1996–2008, (2022).
- [40] Al-Salti, Nasser and Karimov, Erkinjon and Sadarangani, Kishin; On a differential equation with Caputo-Fabrizio fractional derivative of order $1 \leq \beta \leq 2$ and application to mass-spring-damper system, *arXiv preprint arXiv:1605.07381*, (2016).
- [41] Tagne, AS and Ema'a Ema'a, JM and Ben-Bolie, GH and Buske, D; A new truncated M-fractional derivative for air pollutant dispersion, *Indian Journal of Physics*, 94, 11, 1777–1784, (2020).
- [42] Zafar, Asim, Ahmet Bekir, M. Raheel, and Waseem Razzaq. "Optical soliton solutions to Biswas–Arshed model with truncated M-fractional derivative." *Optik* 222: 165355,(2020).
- [43] Jena, Rajarama Mohan and Chakraverty, Snehashish and Baleanu, Dumitru; A novel analytical technique for the solution of time-fractional Ivancevic option pricing model, *Physica A: Statistical Mechanics and its Applications*, 550, 124380, (2020).
- [44] Chen, Qiliang and Baskonus, Haci Mehmet and Gao, Wei and Ilhan, Esin; Soliton theory and modulation instability analysis: The Ivancevic option pricing model in economy, *Alexandria Engineering Journal*, 61, 10, 7843–7851, (2022).
- [45] Chen, Yu-Qiong and Tang, Yu-Hao and Manafian, Jalil and Rezazadeh, Hadi and Osman, MS; Dark wave, rogue wave and perturbation solutions of Ivancevic option pricing model, *Nonlinear Dynamics*, 105, 3, 2539–2548, (2021).