

**Table 1. Demographic and operative details of the case series participants.**

Case	Age (years)	Parity	Gestational age	Previous CS	Antenatal diagnosis with intraoperative confirmation	Bladder displacement in (min)	IPL occlusion (min)	placental separation (min)	ElNoury compression suture (Sec)	EBL (ml)	pRBC (Units)	Postoperative hospital stay (days)
1	36	G2P1	37	1	Accreta + PP	20	10	5	60	800	0	2
2	35	G2P1	39	1	Accreta + PP	10	18	8	60	1000	0	2
3	34	G6P5	36	5	Accreta + PP	12	12	6	60	1400	2	2
4	35	G3P2	36	2	Accreta + PP	9	10	5	60	1300	2	2
5	34	G3P2	36	2	Accreta + PP	20	8	3	60	1400	3	1
6	30	G5P3	37	1	Accreta + PP	6	12	2	120	900	0	1
7	24	G2P1	38	1	PP	7	6	3	60	700	0	1
8	30	G6P4	37	4	Percreta + PP	8	10	6	120	1400	5	3
9	36	G3P2	36	2	Accreta + PP	6	6	4	60	700	0	2
10	38	G4P3	36	3	Increta + PP	3	18	5	60	750	0	3
11	38	G5P4	36	4	Accreta + PP	15	15	13	80	750	0	1
12	34	G4P1	36	1	Increta + PP	8	14	6	80	2000	3	2
13	31	G3P2	36	2	Percreta + PP	20	12	3	80	800	0	2
14	36	G5P4	36	4	Accreta + PP	6	18	5	60	2500	4	3
15	30	G3P2	37	2	Accreta + PP	5	9	2	50	2000	4	2
16	31	G2P1	37	1	Accreta + PP	5	15	4	70	800	0	2

\*No: Case number; \*GA: gestational age (week); \*G/P: Gravidity/Parity; \*CS: number of previous cesarean section; \*PP: placenta praevia; \*IPL: infundibulopelvic ligament; \*EBL: estimated blood loss; \*pRBC: transfused units of packed red blood cells intraoperative and postoperative