

TABLE 3. Genetic patterns of *T. gondii* isolated from chickens classified by continent.

Continent	Country / region* (No. of articles)	No. of isolates or strains	No. of Type (%)			No. of atypical (%)			No. of Not defined (%)	No. of genotypes
			Type I / #10	Type II / #1	Type III / #2	Atypical	Mixed			
America	Brazil (16)	253	1 (0,4%)	5 (2,0%)	1 (0,4%)	142 (56,1%)	1 (0,4%)	103 (40,7%)	105	
	Nicaragua (1)	48	-	-	-	-	-	48 (100,0%)	8	
	Guyana (1)	36	-	-	-	-	-	36(100,0%)	9	
	USA (2)	35	-	11 (31,4%)	12 (34,3%)	12 (34,3%)	-	-	15	
	Argentina (2)	39	2 (5,2%)	-	4 (10,2%)	33 (84,6%)	-	-	6	
	Costa Rica (1)	32	-	-	-	-	-	32 (100,0%)	5	
	Chile (1)	22	-	-	-	-	-	22 (100,0%)	3	
	Saint Kitts (1)	21	-	6 (28,6%)	1 (4,8%)	14 (66,6%)	-	-	6	
	Grenada (1)	20	-	-	15 (75,0%)	5 (25,0%)	-	-	4	
Africa	Caribbean* (1)	11	-	-	2 (18,2%)	4 (36,4%)	-	5 (45,4%)	5	
	<b>Subtotal (27)</b>	<b>517</b>	<b>3 (0,6%)</b>	<b>22 (4,3%)</b>	<b>35 (6,8%)</b>	<b>210 (40,6%)</b>	<b>1 (0,2%)</b>	<b>246 (47,5%)</b>	<b>166</b>	
	Central and Northern Africa* (1)	21	-	-	-	-	-	21 (100,0%)	19	
	Uganda (1)	20	1 (5,0%)	8 (40,0%)	6 (30,0%)	-	5 (25,0%)	-	4	
Asia	<b>Subtotal (2)</b>	<b>41</b>	<b>1 (2,5%)</b>	<b>8 (19,5%)</b>	<b>6 (14,6%)</b>	<b>0 (0,0%)</b>	<b>5 (12,2%)</b>	<b>21 (51,2%)</b>	<b>23</b>	
	China (1)	3	-	-	-	3 (100,0%)	-	-	1	
	<b>Subtotal (1)</b>	<b>3</b>	<b>0 (0,0%)</b>	<b>0 (0,0%)</b>	<b>0 (0,0%)</b>	<b>3 (100,0%)</b>	<b>0 (0,0%)</b>	<b>0 (0,0%)</b>	<b>1</b>	
<b>TOTAL</b>	<b>30</b>	<b>561</b>	<b>4 (0,8%)</b>	<b>30 (5,3%)</b>	<b>41 (7,3%)</b>	<b>213 (37,9%)</b>	<b>6 (1,1%)</b>	<b>267 (47,6%)</b>	<b>190</b>	

\*The study gathered genotypes from more than one country for the article = regional study. “-“= There is not data. “No”= Number.