

TABLE 2 Incidence of GVHD by donor relation, stem cell source, and HLA-match

	Related donor bone marrow or peripheral blood				Unrelated donor bone marrow or Peripheral blood				Cord blood (related or unrelated)			
GVHD	MRD		Haplo M		≥ 8/8 Ag M		≤7/8 Ag M		6/6 Ag M		<6/6 Ag M	
	N	%	N	%	N	%	N	%	N	%	N	%
No aGVHD	50	70.4	14	63.6	21	58.3	5	31.3	9	69.2	7	36.8
Grade I	11	15.5	4	18.2	9	25.0	2	12.5	2	15.4	4	21.1
Grade II	4	5.6	4	18.2	5	13.9	1	6.3	2	15.4	6	31.6
Grade III	4	5.6	0	0	0	0	3	18.8	0	0	0	0
Grade IV	2	2.8	0	0	1	2.8	5	31.3	0	0	2	10.5
Total	71	100	22	100	36	100	16	100	13	100	19	100
No cGVHD	58	81.7	19	86.4	26	72.2	12	75	11	84.6	14	73.7
Limited cGVHD	5	7.0	1	4.5	4	11.1	0	0	2	15.4	4	21.1
Extensive cGVHD	8	11.3	2	9.1	6	16.7	4	25	0	0	1	5.3
Total	71	100	22	100	36	100	16	100	13	100	19	100

HLA data were missing on 6 patients, thus the table includes 177 patients

MRD= matched related donor

Ag M = antigen match